


FIGURE 1

...

ERß antigen


	EBR 14CB (Int 904)	A (Int 90.		FRR 14C8 (lot 002)	3 (lot 00		ERa 1F3 (lot 904)	ot 904)	ш	ERa PAb (lot 904)	(lot 904)		Normal Serum	E	
									l						
Dilution											1		(ć	•
1/250	0.773	0.726 0	0.7495	0.728	0.726	0.727	0	0	0	0.129	0.065 0.097	0.097	0	>	-
1/500	0.513	0.469	0.491	0.571	0.485	0.528	0	0	0	0.028	0.005	0.0165	0	0	0
1/1000	0.303	0.256	_	0.339	0.302	0.3205	0	0	0	0	0	0	0	0	0
000077	2 6	980	0.086 0.1075		0 117	0.13	c	c	0	0	0	0	0	0	0
1/4000		0.000	0.000 0.1075		0.004	0.0115	0	0	-	0	0	0	0	0	



·:. .


$\mathsf{ER}\alpha$ antigen

	ERB 14C8 (lot 904)	8 (lot 90		ERB 14C8 (lot 002)	8 (lot 00		ERa 1F3 (lot 904)	(lot 904)		ERa PAb (lot 904)	(lot 904		Normal Serum	erum	
Dilution															
1/250	0.017	0.015	0.015 0.016	0	0	0	0.739	0.858	0.7985	2.681	2.73	2.7055	0.047	0.055	0.051
1/500	0	0.004	0.002	0.003	0.007	0.005	0.539	0.394	_	1.962	1.996	1.979	0.022	0.023	0.0225
1/1000	0.009	0	0.0045	9000	0.008	0.007	0.287	0.279	0.283		1.568	1.5055	0.009	0.025	0.017
1/2000	0	0.012	0.012 0.006	0	0.009	0.0045	0.278	0.12			0.888	0.929	0	0	0
1/4000	0.012	0.012 0.005 0.0085	0.0085	0.016	0	0.008	0.061	0.081	0.071	0.617	0.505	0.561	0	0	0



••

Immunocytochemical analysis of paraffin-embedded, fixed tissue processed with 14C8 ER-ß monoclonal antibody.



B. Infiltrating lobular carcinoma of the breast.